

Use of a standardized runoff index for characterizing hydrologic drought

Andrew W. Wood¹ and Shraddhanand Shukla¹

¹Department of Civil and Environmental Engineering, Box 352700, University of Washington,
Seattle, WA 98115

1 **Abstract**

2 Many current metrics of drought are derived solely from analyses of climate variables such
3 as precipitation and temperature. Drought is clearly a consequence of climate anomalies, as well
4 as of human water use practices, but many impacts to society are more directly related to
5 hydrologic conditions resulting from these two factors. Modern hydrology models can provide a
6 valuable counterpart to existing climate-based drought indices by simulating variables such as
7 land surface runoff. We contrast the behavior of a standardized runoff index (SRI) with that of
8 the well-known standardized precipitation index (SPI) during drought events in a snowmelt
9 region. Although the SRI and SPI are similar when based on long accumulation periods, the SRI
10 incorporates hydrologic processes that determine seasonal lags in the influence of climate on
11 streamflow. As a result, on monthly to seasonal time scales, the SRI is a useful counterpart to
12 the SPI for depicting hydrologic aspects of drought.

1 **1. Motivation**

2 Drought is a multi-faceted phenomenon that occurs across a range of temporal and spatial
3 scales and is experienced across a range of societal sectors that are dependent on climate and
4 water resources (Wilhite, 2000). Many indicators used to describe drought are derived from
5 analyses of climate variables such as precipitation and temperature. The impacts of drought are a
6 consequence of climate anomalies, as well as of human water use practices, but many impacts
7 are more directly related to the resulting hydrologic conditions. Several drought indices attempt
8 to incorporate the interaction of the land surface with climate, including, e.g., the Palmer
9 Drought Severity Index (PDSI; Palmer, 1965), but the conceptual moisture accounting of the
10 PDSI (and related indices) represents evaporative effects crudely and omits snow accumulation
11 and melt entirely (Allen, 1984). Others, such as the Standardized Precipitation Index (SPI;
12 McKee et al., 1993) are purely related to climate. Soil moisture is also recognized as an
13 important indicator of drought for agriculture and other sectors. The National Oceanic and
14 Atmospheric Administration (NOAA) Climate Prediction center (CPC), for example, uses a
15 simplified water balance model (Huang et al. 1996) to simulate soil moisture percentiles for
16 drought monitoring in the United States (US). The current state-of-the-practice drought analysis
17 in the US, the US Drought Monitor (USDM; Svoboda et al., 2002), is subjectively assembled
18 from estimates of climate indices such as the such as the PDSI, SPI, streamflow percentiles,
19 snow water equivalent (SWE) anomalies, CPC soil moisture percentiles, and various other minor
20 inputs.

21 The potential to advance the hydrologic basis for drought monitoring now exists. The last
22 decade has seen the maturation of a class of macro-scale hydrology models that incorporate
23 sufficient physics to maintain water and energy balances of the major components of the top 1-2

1 meters of the land surface, operate at between an hourly and daily timestep, and run at horizontal
2 grid resolutions that are far finer than the climate division and even county level data sources of
3 the USDM. Foremost among the modeling efforts in the US is the NOAA Land Data
4 Assimilation Project (NLDAS; Mitchell et al., 2003), which has helped to promote the
5 development and calibration of four land surface schemes over much of North America at 1/8
6 degree spatial resolution.

7 These models produce real-time modeled soil moisture, SWE and runoff estimates at sub-
8 daily time steps. We focus here on runoff, a primary concern to water managers, because it is
9 closer to being a verified product from models than soil moisture. The drought monitoring and
10 management community widely uses data expressed in an integer-based index framework. We
11 also employ an index framework to demonstrate the application of modeled runoff for water
12 cycle analysis in the context of drought.

13 **2. Hydrologic Modeling**

14 We use the physically based, semi-distributed macro-scale Variable Infiltration Capacity
15 (VIC) model (Liang et al. 1994) to simulate the land surface water balance. Like other NLDAS
16 project models, VIC accounts for modulation by vegetation of land-atmosphere moisture and
17 energy fluxes, and attempts to represent sub-grid variability of vegetation, soil and terrain
18 characteristics via sub-grid area-specific parameter classifications and an infiltration algorithm
19 that involves the areal extent of soil saturation. Soil layer depths, infiltration and base flow
20 parameters are adjusted during model calibration, the primary goal of which is often to reproduce
21 observed streamflow.

22 The VIC model has been applied for analyzing historical drought events for the entire U.S.
23 (Andreadis et al., 2005) and globally (Sheffield and Wood, 2007). We draw here on VIC model

1 results from the 1/8th degree implementation for the Feather R. basin, California, described in
2 Wood and Schaake (2007), and on the 1/2 degree continental US (CONUS) implementation
3 (Andreadis et al, 2005) that is incorporated in the real-time University of Washington Surface
4 Water Monitor (see <http://www.hydro.washington.edu/forecast/monitor/>). Both models use
5 gridded NOAA National Climatic Data Center Cooperator station precipitation, temperature
6 minima and maxima as inputs to simulate the water balance at a daily timestep. Typical outputs
7 include daily surface runoff, baseflow, evaporation, soil moisture and SWE for each grid cell.
8 We sum daily surface runoff and baseflow from each cell together to form, simply, “runoff”, and
9 temporally average runoff to form monthly totals for each model grid cell. At large spatial
10 scales, this runoff differs from streamflow because it is not routed through a channel network.

11 **3. Standardized Runoff Index**

12 We apply the concept employed by Mckee et al. (1993) for the SPI in defining a standardized
13 runoff index (SRI) as the unit standard normal deviate associated with the percentile of
14 hydrologic runoff accumulated over a specific duration. Different durations (e.g., 1-month, 9-
15 month) and different spatial aggregations of the index can be calculated depending on source
16 data resolution and desired application. SPIs, for instance, are calculated by NOAA on a climate
17 division basis and by state agencies on a county level basis. The procedure for calculating the
18 SRI includes the following steps:

- 19 1. A retrospective time series of runoff is obtained by simulation, and a probability
20 distribution is fit to the sample represented by the time series values.
- 21 2. The distribution is used to estimate the cumulative probability of the runoff value of
22 interest (either the current accumulation or one from a retrospective date).

1 3. The cumulative probability is converted to a standard normal deviate (with zero mean
2 and unit variance), which can either be calculated from a numerical approximation to the
3 normal cumulative distribution function (CDF) or extracted from a table of values for the
4 normal CDF that is readily available in statistics textbooks or on the World Wide Web.

5 McKee et al. (1993) select the Gamma distribution for fitting monthly precipitation data
6 series, and suggest that the procedure can be applied to other variables relevant to drought, e.g.,
7 streamflow or reservoir contents. In pursuing this suggestion for model-based runoff, we note
8 that distributions other than the Gamma may be more appropriate, depending on the runoff
9 variable's retrospective sample characteristics (especially skew and kurtosis), which vary greatly
10 by geographic location and degree of temporal aggregation. **Figure 1**, showing the distribution
11 of areal average 3-month total simulated runoff during March, April and May from the years
12 1955-2005 for the Feather River basin, California, makes clear that any value of runoff can
13 equally well be expressed in terms of its percentile (top axis) or the standardized index (bottom
14 axis). Here the 2 parameter log normal (LN) distribution, for example, provides a better fit at
15 high extremes than the Gamma distribution whereas the Gamma distribution may perform better
16 for low runoff values. The 3-parameter LN and Generalized Extreme Value distributions may
17 have even better general applicability for runoff over widely varying hydro-climatic regimes.
18 Where a satisfactory distribution fit cannot be achieved, an alternative is to estimate percentiles
19 empirically. Care must be taken to negotiate the first two steps above so as to minimize errors in
20 estimating the probability of runoff, particularly in arid regions where runoff may be
21 intermittent. Note that SRI units are non-linearly related to percentiles, so a change in SRI from -
22 2 to -3 equals a smaller percentile change than an SRI change from 0 to 1. This correspondence

1 is common to other USDM indices, reflecting the non-linear relationship of drought impacts to
2 the probability of a drought event.

3 **4. Results**

4 We illustrate differences in the behavior of the SPI and SRI using areal averages of observed
5 precipitation and simulated runoff in the Feather River basin. **Figure 2** shows monthly time
6 series of the indices for four accumulation periods (1-, 3-, 6- and 12-month) for the years 1975-
7 1995. The 12-month SPI and SRI are very similar due to the high correlation between annual
8 precipitation and runoff ($r=0.9$). The differences between the indices increase as the
9 accumulation period decreases, with SPI to SRI correlations dropping from 0.88 for the 12-
10 month period to 0.82, 0.69 and 0.01 for the 6-, 3-, and 1-month period indices. For shorter
11 period accumulations, the SRI is less variable from month to month than the SPI due to the
12 detention of moisture in snow or soil storages that regulate runoff . Supplemental Figure S1
13 shows the autocorrelation of 1-month runoff provided by these storages at different times of the
14 year. The 12-month indices integrate over an entire water year, longer than most effects of
15 hydrologic modulation. A long integration, however, accumulates values that are well past and
16 may no longer influence current land surface conditions. For example, a large event in early
17 1986 causes the 12-month indices to remain above 1.0 for 2 months after the 1- and 3-month
18 SRIs fall below values of -1.0, in December, 1986.

19 The 1976-1977 and 1987-1992 California drought events are evident in all four sets of
20 timeseries, as is the 1995 flood year. During the drought years, the 12-month indices are both
21 negative, with the exception of a few months in 1989. These results compare well qualitatively
22 to recorded precipitation and runoff anomalies for Sacramento River basin, which contains the
23 Feather River basin: percentages of normal during each of the six years were, respectively, 55,

1 75, 100, 75, 75, 76 for precipitation and 49, 49, 78, 49, 45, 47 for runoff (Dziegielewski et al,
2 1993). During drought events, the shorter period indices attain positive values more frequently
3 than the 12-month indices and the SPI recovers to above normal levels more frequently than the
4 SRI. In 1990-1991, the SPI describes spells of precipitation that are insufficient to ameliorate
5 hydrologic drought conditions, a reality reflected in the non-recovery of the SRI at these times.
6 In winter 1992-1993, however, the SPI increase corresponds to a building snowpack, and
7 presages a recovery from drought; whereas the SRI reflects only current runoff and remains low
8 until the snowmelt period arrives. In this case, the SPI contains predictive information not
9 present in the SRI.

10 The hydrologic importance of precipitation varies greatly depending on the seasonal
11 precipitation climatology and the current land surface moisture state. **Figure 3** shows the 3-
12 month SPI and SRI values for the Feather R. basin together with daily basin areal averages of
13 observed accumulated water year precipitation and simulated soil moisture, SWE and water year
14 runoff for the drought year of 1991 and the flood year of 1995. In 1991, recorded precipitation
15 in March was 300 percent of average, which is evident in the step change in accumulated
16 precipitation, soil moisture and runoff, increases in SWE, and return to above normal of the 3-
17 month SPI. The recharge of low soil moisture, however, diminishes runoff, muting the SRI
18 recovery. Qualitatively, these dynamics are corroborated by reports of Dziegielewski et al
19 (1993): “Since the onset of drought in 1987, California has experienced at least one month of
20 above-normal precipitation during each water year...However, these precipitation ‘bursts’ were
21 not adequate to overcome water shortages in most parts of the state accumulated during the
22 previous months of the respective water years.”

1 A hydrologic contrast is given by the extreme precipitation events in January and March
2 1995, which together caused flooding that claimed 28 lives and led to over \$220 million in
3 damages (California, 2003). As Figure 3 shows, these storms arrived when soil moisture was
4 near or above normal, and drove increases in SWE, runoff and consequently both SPI and SRI to
5 much above normal. Below average summer precipitation then caused SPI to plummet.
6 Precipitation during California’s relatively dry summers, however, often contributes less to
7 summer runoff than the moisture accumulated in the soil and SWE during winter. In 1995, the
8 summer precipitation deficits were greatly overshadowed by the discharge of stored moisture,
9 hence runoff and the SRI remained high through the end of the water year.

10 Due to hydrologic delays in the form of snow and soil moisture, the SPI may be
11 desynchronized from response of the land surface to those anomalies (the USDM, partly for this
12 reason, advises caution in interpreting short duration SPIs) – whereas the SRI incorporates the
13 land surface dynamics that moderate the hydrologic response. Snow-free regions lack the delay
14 due to snow accumulation (longer than soil storage delay), hence the SPI and SRI are better
15 correlated at short durations, despite the SRI’s attenuation of the precipitation signal.
16 Supplemental Figures S2 and S3 duplicate Figures 2 and 3 for a snow-free location.

17 **5. Operational mapping of runoff on continental scales**

18 The land surface modeling activities noted in section 1 are capable of producing runoff and
19 derived indices such as the SRI on a continental scale, using multiple hydrologic models, for
20 both current and projected conditions (e.g., based on climate predictions from CPC). **Figure 4**,
21 for example, shows a real-time SRI analysis based on the ½ degree VIC model simulation
22 (Section 2) for 3-month runoff conditions as of September 2007. In general, the locations of wet
23 and dry conditions across the domain are represented similarly by the SPI and SRI. The SRI is

1 higher than the SPI in Oklahoma and Texas due to surplus moisture remaining from extreme
2 summer 2007 flooding, and is lower in the western US where the water year was much drier than
3 average. In California, the 3-month SPI is above normal, in contrast to the SRI and to the 2007-
4 09-25 USDM, which depicted all counties as being at least “Abnormally Dry”. The recent
5 extension of NLDAS retrospective simulations back to 1979 is long enough estimate runoff
6 distributions, despite post-dating important prior drought events (e.g., 1976-77, the Dust Bowl).
7 The potential value of hydrologic anomaly products that depend on the existence of consistent
8 historical analyses argues that the NLDAS extension should be a stepping stone toward even
9 longer retrospective simulations.

10 **6. Conclusions**

11 The foregoing examples demonstrate that modeled runoff could provide a useful counterpart
12 to climate-based indices for drought monitoring and management. Due to the popularity of the
13 SPI, the multi-period SRI framework for runoff is likely to be familiar to the drought research,
14 monitoring and management communities. Whereas the SPI describes the climate anomalies in
15 isolation from their hydrologic context, the SRI directly describes the effects of climate
16 anomalies on current hydrologic conditions as governed by land surface physical processes. One
17 strength of a runoff-based index is that it can be forecast, and its predictability depends not only
18 on climate outlooks, for which seasonal skill is generally low, but on hydrologic initial
19 conditions, which in some seasons largely determine future runoff (e.g., spring snow state in the
20 western US). A second strength is that calibrated runoff simulations are more widely available
21 for real-time application than naturalized (e.g., adjusted to remove human impairments)
22 streamflow observations, which precludes the use of streamflow in a real-time framework (e.g.,
23 Modarres, 2007) in most locations. A weakness of the SRI is that it is based on modeled runoff

1 that cannot be verified everywhere, and it reflects the customary uncertainties associated with
2 models. Nevertheless, model-based hydrologic runoff (in index form or otherwise) shows
3 potential to complement existing climate indices and local hydro-climatological information
4 (e.g., knowledge of when/where snow accumulation is occurring), leading to improvements in
5 the assessment of current and future drought status.

6 .

7 **Acknowledgements**

8 The authors are grateful for the advice of Steve Burges; to Doug Lecomte, Victor Murphy
9 and Mark Svoboda for feedback on hydrologic modeling in drought monitoring; and for the
10 funding support from the NOAA TRACS program under Cooperative Agreement
11 NAXXXXXXXX.

12 **References**

- 13 Alley, W. M. (1984), The Palmer Drought Severity Index: Limitations and assumptions, *J. Clim.*
14 *Appl. Meteorol.*, 23, 1100– 1109.
- 15 Andreadis, K.M., E.A. Clark, A.W. Wood, A.F. Hamlet, and D.P. Lettenmaier (2005), 20th
16 Century Drought in the Conterminous United States, *J. Hydrometeor.*, 6(6): 985-1001.
- 17 California (2003), High water in California. State of California, the Resources Agency, Dept. of
18 Water Resources.
- 19 Dziegielewski, Benedykt, Hari P. Garbharran, and John F. Langowski (1993), Lessons learned
20 from the California drought (1987-1992). IWR report, 93-NDS-5. Fort Belvoir, Va: The
21 Institute.

1 Huang, J., H. M. Van den Dool, and K. P. Georgakakos (1996), Analysis of model-calculated
2 soil moisture over the United States (1931-1999) and applications to long-range
3 temperature forecasts. *J. Climate*, 9, 1350-1362.

4 Liang, X., D. P. Lettenmaier, E. F. Wood, and S. J. Burges (1994), Simple hydrologically Based
5 Model of Land Surface Water and Energy Fluxes for GSMs, *J. Geophys. Res.*, **99**(D7),
6 14,415-14,428.

7 McKee, T. B., N. J. Doesken, and J. Kleist, (1993), The relationship of drought frequency and
8 duration to timescales, paper presented at 8th Conference on Applied Climatology,
9 Anaheim, Calif., 17– 22 Jan.

10 Mitchell, K. E., and many authors (2004), The multi-institution North American Land Data
11 Assimilation System (NLDAS): Utilizing multiple GCIP products and partners in a
12 continental distributed hydrological modeling system, *J. Geophys. Res.*, 109, D07S90,
13 doi:10.1029/2003JD003823.

14 Modarres, R., (2007), Streamflow drought time series forecasting, *Stoch Environ Res Risk Assess*
15 21:223–233, DOI 10.1007/s00477-006-0058-1

16 Palmer, W. C. (1965), Meteorological drought, Res. Pap. 45, 58 pp., *Weather Bur.*, U.S. Dep. of
17 Commer., Washington, D. C.

18 Sheffield, J., and E. F. Wood, 2007, Global trends and variability in soil moisture and drought
19 characteristics, 1950-2000, from observation driven simulations of the terrestrial
20 hydrologic cycle. *J. Climate*, (in press).

21 Svoboda, M., and many authors, 2002: The Drought Monitor. *Bull. Amer. Meteor. Soc.*, **83**,
22 1181–1190.

- 1 Wilhite, D. A. (2000), Drought as a natural hazard: Concepts and definitions, in Drought: A
- 2 Global Assessment, Hazards Disasters Ser., vol. I, edited by D. A. Wilhite, pp. 3– 18,
- 3 Routledge, New York.
- 4 Wood, A.W. and J.C. Schaake, 2007, Correcting errors in streamflow forecast ensemble mean
- 5 and spread, *J. Hydrometeorology* (in press).

1 **FIGURE CAPTIONS**

2 **Figure 1** The simulated historical distribution of total runoff (areal average depth)
3 accumulated over March, April and May for the Feather River basin for the period
4 1955-2005. The sample is fitted with Gamma and Log Normal (LN2)
5 distributions. The SRI (bottom axis) is the unit standard normal deviate
6 associated with the percentiles of the runoff value (top axis).

7 **Figure 2** Historical time series of the SPI and SRI for 1-, 3-, 6- and 12-month accumulation
8 periods, based on observed precipitation and simulated runoff in the Feather River
9 basin.

10 **Figure 3** The Feather River basin areal average water balance during two water years (1991
11 left; 1995 right) compared with the 3-month SPI and SRI. Daily values of
12 observed precipitation and simulated soil moisture, snow water equivalent and
13 runoff in each of the two years (red line) are plotted against the minimum,
14 maximum and quartiles of their daily historical distribution from 1955-2005. The
15 SPI and SRI have a monthly timestep.

16 **Figure 4** The 3-month SPI and SRI for September 2007 as calculated from observed
17 precipitation and simulated runoff on a ½ degree grid.

FIGURES




Figure 1

SPI and SRI for the Feather R. Basin, CA




Figure 2

Feather River Basin at Oroville, CA
 (current (red) 1955–2005 min, max and quartiles)





Figure 3

Standardized Precipitation Index (3-Month) 200709



Standardized Runoff Index (3-Month) 200709




Figure 4